

# Extended formulations in combinatorial optimization: constructions and lower bounds

Samuel Fiorini

Université libre de Bruxelles (ULB, Brussels)

VI IPCO Summer School

VIII Escuela de Verano en Matemáticas Discretas

Day 1

# Today

- 1 Introduction
- 2 Basics on polytopes and polyhedra
- 3 Factorization theorem


# Today

- 1 Introduction
- 2 Basics on polytopes and polyhedra
- 3 Factorization theorem

# Linear programming

## Example


$$\begin{array}{ll}\max & 4x_1 + 3x_2 \\ \text{subject to} & -x_1 + 4x_2 \leq 16 \\ & x_1 + x_2 \leq 9 \\ & 3x_1 - x_2 \leq 15 \\ & x_1 \geq 0 \\ & x_2 \geq 0\end{array}$$



# Linear programming

## Example

$$\begin{array}{ll}\max & 4x_1 + 3x_2 \\ \text{subject to} & -x_1 + 4x_2 \leq 16 \\ & x_1 + x_2 \leq 9 \\ & 3x_1 - x_2 \leq 15 \\ & x_1 \geq 0 \\ & x_2 \geq 0\end{array}$$



**LP in general form:**

$$\begin{array}{ll}\max & \sum_{i=1}^d w_i x_i = w^\top x \\ \text{subject to} & Ax \leq b\end{array}$$

# Linear programming



A successful tool

## Fact

Linear programming is **very successful** both in practice and theory

3 reasons for this success:

- 1 Many **problems** can be expressed as LPs






- 2 Rich **theory**: LP duality can certify the quality of solutions

$$\begin{array}{ll} \max & w^T x \\ \text{s.t.} & Ax \leq b \end{array} = \begin{array}{ll} \min & b^T y \\ \text{s.t.} & A^T y = w \\ & y \geq 0 \end{array}$$

- 3 There exist powerful **algorithms** for solving LPs




# Linear programming

## Algorithms

Algorithm / method	Poly-time?	In practice?	Generalizes to convex?
 (simplex)	No	<b>Very fast</b>	No
 (ellipsoid)	<b>Yes</b>	Slow	Yes
 (int. point)	Yes	Fast	<b>Yes</b>

# Linear programming

## Algorithms




Algorithm / method	Poly-time?	In practice?	Generalizes to convex?
 (simplex)	No	<b>Very fast</b>	No
 (ellipsoid)	<b>Yes</b>	Slow	Yes
 (int. point)	Yes	Fast	<b>Yes</b>

All practical algorithms have sensitivity to **size** of input LP



# Linear programming

## Algorithms


Algorithm / method	Poly-time?	In practice?	Generalizes to convex?
 (simplex)	No	<b>Very fast</b>	No
 (ellipsoid)	<b>Yes</b>	Slow	Yes
 (int. point)	Yes	Fast	<b>Yes</b>

All practical algorithms have sensitivity to **size** of input LP

⇒ try to minimize size: variables, constraints are **resources**

# Combinatorial optimization


## LP formulation of TSP



**TSP:** Given a graph  $G = (V, E)$  and distances  $d_{ij}$  for each  $ij \in E$ , find tour of minimum length.

# Combinatorial optimization


## LP formulation of TSP




**TSP:** Given a graph  $G = (V, E)$  and distances  $d_{ij}$  for each  $ij \in E$ , find tour of minimum length.

$$\min \sum_{ij \in E} d_{ij} x_{ij}$$


s.t.  $x \in \{0, 1\}^{|E|}$  encodes tour of  $G$



$(1, 0, 1, 1, 0, 1)$



$(1, 1, 0, 0, 1, 1)$




$(0, 1, 1, 1, 1, 0)$

$\mathbb{R}^6$



# Combinatorial optimization


## LP formulation of TSP




**TSP:** Given a graph  $G = (V, E)$  and distances  $d_{ij}$  for each  $ij \in E$ , find tour of minimum length.

$$\min \sum_{ij \in E} d_{ij} x_{ij}$$


$$\text{s.t. } x \in \{0, 1\}^{|E|} \text{ encodes tour of } G$$
$$x \in \text{TSP}(G)$$



$$(1, 0, 1, 1, 0, 1)$$




$$(1, 1, 0, 0, 1, 1)$$



$$(0, 1, 1, 1, 1, 0)$$

$\mathbb{R}^6$




# Combinatorial optimization

## TSP polytope

TSP polytope of  $G$ :

$$\text{TSP}(G) := \text{conv}\{x \in \{0,1\}^{|E|} \mid x \text{ encodes tour of } G\}$$




# Combinatorial optimization

## TSP polytope

TSP polytope of  $G$ :

$$\text{TSP}(G) := \text{conv}\{x \in \{0,1\}^{|E|} \mid x \text{ encodes tour of } G\}$$




$$\begin{array}{ll} \min & \sum_{ij \in E} d_{ij} x_{ij} \\ \text{s.t.} & x \in \text{TSP}(G) \end{array}$$

# Combinatorial optimization

## TSP polytope

TSP polytope of  $G$ :

$$\text{TSP}(G) := \text{conv}\{x \in \{0,1\}^{|E|} \mid x \text{ encodes tour of } G\}$$




$$\begin{array}{ll} \min & \sum_{ij \in E} d_{ij} x_{ij} \\ \text{s.t.} & x \in \text{TSP}(G) \\ & Ax \geq b \end{array}$$

# Combinatorial optimization

## TSP polytope

TSP polytope of  $G$ :

$$\text{TSP}(G) := \text{conv}\{x \in \{0,1\}^{|E|} \mid x \text{ encodes tour of } G\}$$



$$\begin{array}{ll} \min & \sum_{ij \in E} d_{ij} x_{ij} \\ \text{s.t.} & \cancel{x \in \text{TSP}(G)} \\ & Ax \geq b \end{array}$$

**Issue:** System  $Ax \geq b$  for  $\text{TSP}(G)$  is




# Combinatorial optimization

## TSP polytope

TSP polytope of  $G$ :

$$\text{TSP}(G) := \text{conv}\{x \in \{0,1\}^{|E|} \mid x \text{ encodes tour of } G\}$$



$$\begin{array}{ll} \min & \sum_{ij \in E} d_{ij} x_{ij} \\ \text{s.t.} & \cancel{x \in \text{TSP}(G)} \\ & Ax \geq b \end{array}$$


**Issue:** System  $Ax \geq b$  for  $\text{TSP}(G)$  is huge,

# Combinatorial optimization

## TSP polytope

TSP polytope of  $G$ :

$$\text{TSP}(G) := \text{conv}\{x \in \{0,1\}^{|E|} \mid x \text{ encodes tour of } G\}$$



$$\begin{array}{ll} \min & \sum_{ij \in E} d_{ij} x_{ij} \\ \text{s.t.} & \cancel{x} \in \cancel{\text{TSP}(G)} \\ & Ax \geq b \end{array}$$

**Issue:** System  $Ax \geq b$  for  $\text{TSP}(G)$  is huge, not known

# Extended formulations

## Reformulating an LP

In general, we hope to go from this:

$$\min \sum_{i=1}^d w_i x_i$$

$$\text{s.t. } Ax \geq b$$

← large, not known, complicated

# Extended formulations

## Reformulating an LP

In general, we hope to go from this:

$$\begin{array}{ll} \min & \sum_{i=1}^d w_i x_i \\ \text{s.t.} & Ax \geq b \end{array} \quad \leftarrow \text{large, not known, complicated}$$

...to this:

$$\begin{array}{ll} \min & \sum_{i=1}^d w_i x_i \\ \text{s.t.} & Ex + Fy = g, y \geq \mathbf{0} \end{array} \quad \leftarrow \text{small, explicit, simple}$$

# Extended formulations

## Reformulating an LP

In general, we hope to go from this:

$$\begin{array}{ll} \min & \sum_{i=1}^d w_i x_i \\ \text{s.t.} & Ax \geq b \end{array} \quad \leftarrow \text{large, not known, complicated}$$

... to this:

$$\begin{array}{ll} \min & \sum_{i=1}^d w_i x_i \\ \text{s.t.} & Ex + Fy = g, y \geq 0 \end{array} \quad \leftarrow \text{small, explicit, simple}$$

In such a way that:  $Ax \geq b \iff \exists y : Ex + Fy = g, y \geq 0$

# Extended formulations

## Reformulating an LP

In general, we hope to go from this:

$$\begin{array}{ll} \min & \sum_{i=1}^d w_i x_i f(x) \\ \text{s.t.} & Ax \geq b \end{array} \quad \leftarrow \text{large, not known, complicated}$$

... to this:

$$\begin{array}{ll} \min & \sum_{i=1}^d w_i x_i f(x) \\ \text{s.t.} & Ex + Fy = g, y \geq 0 \end{array} \quad \leftarrow \text{small, explicit, simple}$$

In such a way that:  $Ax \geq b \iff \exists y : Ex + Fy = g, y \geq 0$

# Extended formulations

## Definitions (1/2)

$$P = \{x \in \mathbb{R}^d \mid Ax \geq b\} \text{ polytope in } \mathbb{R}^d$$



# Extended formulations

## Definitions (1/2)

$$P = \{x \in \mathbb{R}^d \mid Ax \geq b\} \text{ polytope in } \mathbb{R}^d$$



Definition (Extended formulation = EF)

$Ex + Fy = g, y \geq \mathbf{0}$  is an **EF** of  $P$  if


$$Ax \geq b \iff \exists y : Ex + Fy = g, y \geq \mathbf{0}$$



# Extended formulations

## Definitions (1/2)

$$P = \{x \in \mathbb{R}^d \mid Ax \geq b\} \text{ polytope in } \mathbb{R}^d$$



## Definition (Extended formulation = EF)

$Ex + Fy = g, y \geq 0$  is an **EF** of  $P$  if

$$Ax \geq b \iff \exists y : Ex + Fy = g, y \geq 0$$

## Definition (Extension)

Polytope  $Q$  in  $\mathbb{R}^e$  is an **extension** of  $P$  if  $\exists$  linear  $\pi$  with  $\pi(Q) = P$

# Extended formulations

## Definitions (2/2)

### Definition (Size)

**Size** of EF  $:=$  #inequalities in EF

**Size** of extension  $:=$  #facets of extension

# Extended formulations

## Definitions (2/2)

### Definition (Size)

**Size** of EF  $:=$  #inequalities in EF

**Size** of extension  $:=$  #facets of extension

### Definition (Extension complexity)

**Extension complexity** of  $P = \text{xc}(P) :=$  minimum size of an EF of  $P$

# Extended formulations

## Definitions (2/2)


### Definition (Size)

**Size** of EF  $:=$  #inequalities in EF

**Size** of extension  $:=$  #facets of extension

### Definition (Extension complexity)

**Extension complexity** of  $P = \text{xc}(P) :=$  minimum size of an EF of  $P$



$$\text{xc}(P) \leq 5 < \text{size}(P)$$

# Extended formulations

## Definitions (2/2)


### Definition (Size)

**Size** of EF  $:=$  #inequalities in EF


**Size** of extension  $:=$  #facets of extension

### Definition (Extension complexity)

**Extension complexity** of  $P = \text{xc}(P) :=$  minimum size of an EF of  $P$



$$\text{xc}(P) \leq 5 < \text{size}(P)$$



$$\text{xc}(P) \leq 6 < \text{size}(P)$$

# Extended formulations

Example: Regular polygons

Theorem (**Ben-Tal & Nemirovski'01**)

*If  $P$  is a regular  $n$ -gon, then  $xc(P) = O(\log n)$  !!!*

# Extended formulations

Example: Regular polygons

Theorem (**Ben-Tal & Nemirovski'01**)

*If  $P$  is a regular  $n$ -gon, then  $xc(P) = O(\log n)$  !!!*

**Exercise.** Prove this using reflections.


# Extended formulations

Example: Regular polygons

Theorem (**Ben-Tal & Nemirovski'01**)

*If  $P$  is a regular  $n$ -gon, then  $xc(P) = O(\log n)$  !!!*

**Exercise.** Prove this using reflections.






# Extended formulations

Example: Regular polygons

Theorem (**Ben-Tal & Nemirovski'01**)

*If  $P$  is a regular  $n$ -gon, then  $xc(P) = O(\log n)$  !!!*

**Exercise.** Prove this using reflections.




# Extended formulations

Example: Regular polygons

Theorem (**Ben-Tal & Nemirovski'01**)

If  $P$  is a regular  $n$ -gon, then  $xc(P) = O(\log n)$  !!!

**Exercise.** Prove this using reflections.




# Extended formulations

Example: Regular polygons

Theorem (**Ben-Tal & Nemirovski'01**)

If  $P$  is a regular  $n$ -gon, then  $xc(P) = O(\log n)$  !!!

**Exercise.** Prove this using reflections.



$$xc(P) = xc(\text{conv}(P_1 \cup \sigma(P_1))) \leq xc(P_1) + 2$$

# Fundamental question and dictionary

Within the framework of EFs, we ask:

*Regarding (variables and) constraints as resources, what problems can be solved via LP with a given amount of resources?*

# Fundamental question and dictionary

Within the framework of EFs, we ask:

*Regarding (variables and) constraints as resources, what problems can be solved via LP with a given amount of resources?*

Dictionary:

Algorithmic	Geometric
Problem	Polytope
Algorithm	EF
Complexity	Size
Hardness result	Lower bound

In this course on extended formulations (EFs), we will see:


- ① many examples of EFs
- ② techniques for constructing EFs
- ③ techniques for proving lower bounds on the size of EFs
- ④ a proof that  $\text{TSP}(K_n)$  has no polynomial-size EF

# Today

- 1 Introduction
- 2 Basics on polytopes and polyhedra
- 3 Factorization theorem

# (Convex) polytopes

- ① A **V-polytope** is the convex hull of a finite set of points



$$P = \text{conv}\{v_1, \dots, v_n\}$$


where

$$v_j \in \mathbb{R}^d \quad \forall j$$



# (Convex) polytopes

- ① A **V-polytope** is the convex hull of a finite set of points




$$P = \text{conv}\{v_1, \dots, v_n\}$$

where

$$v_j \in \mathbb{R}^d \quad \forall j$$

- ② A **H-polytope** is the intersection of a finite # of halfspaces (provided this intersection is bounded)



$$P = \{x \in \mathbb{R}^d \mid h_1(x) \leq 0, \dots, h_m(x) \leq 0\}$$

where

$$h_i(x) = A_i x - b_i \quad \forall i$$

# Fundamental theorem

## Theorem

For all  $P \subseteq \mathbb{R}^d$ :

$P$  is a  $V$ -polytope  $\iff P$  is a  $H$  polytope

**Proof** ( $\implies$ )

$$\begin{aligned} P &= \text{conv}\{v_1, \dots, v_n\} \\ &= \left\{ x \in \mathbb{R}^d \mid \exists y \in \mathbb{R}^n : x = \sum_{j=1}^n y_j v_j, \sum_{j=1}^n y_j = 1, y_j \geq 0 \forall j \right\} \\ &= \left\{ x \in \mathbb{R}^d \mid \exists y \in \mathbb{R}^n : Ex + Fy = g, y \geq 0 \right\} \end{aligned}$$

So  $P$  is the **projection** into  $x$ -space of a  $H$ -polytope! This is our 2nd example of an EF. (**Exercise.** What kind of polytope is  $Q$ ?)

$$P = \text{proj}_x(Q) \quad \text{where} \quad Q = \left\{ (x, y) \in \mathbb{R}^{d+n} \mid Ex + Fy = g, y \geq 0 \right\}$$

# Eliminating one variable (Fourier-Motzkin elimination)

To conclude the proof, it suffices to show that **eliminating** a single variable from a system of linear constraints “produces” a new system of linear constraints (changing notations):

$$\begin{aligned} Q &= \{(x, y) \in \mathbb{R}^{d+1} \mid A_i x + b_i y \leq c_i \ \forall i\} \\ &= \left\{ (x, y) \in \mathbb{R}^{d+1} \mid \begin{array}{lll} A_{i_0} x & \leq & c_{i_0} & \forall i_0 \in I_0 \\ y & \leq & b_{i_+}^{-1} c_{i_+} - b_{i_+}^{-1} A_{i_+} x & \forall i_+ \in I_+ \\ y & \geq & b_{i_-}^{-1} c_{i_-} - b_{i_-}^{-1} A_{i_-} x & \forall i_- \in I_- \end{array} \right\} \end{aligned}$$

where  $I_0 = \{i \mid b_i = 0\}$ ,  $I_+ = \{i \mid b_i > 0\}$ ,  $I_- = \{i \mid b_i < 0\}$

Then  $\text{proj}_x(Q)$  is defined by:

$$\begin{array}{lll} A_{i_0} x & \leq & c_{i_0} & \forall i_0 \in I_0 \\ b_{i_-}^{-1} c_{i_-} - b_{i_-}^{-1} A_{i_-} x & \leq & b_{i_+}^{-1} c_{i_+} - b_{i_+}^{-1} A_{i_+} x & \forall i_+ \in I_+, i_- \in I_- \end{array}$$



# Projections

From proof, see that if  $Q = \{(x, y) \in \mathbb{R}^{d+k} \mid Ax + By \leq c\}$  then

$$\text{proj}_x(Q) = \{x \in \mathbb{R}^d \mid u^\top Ax \leq u^\top c \text{ for finite \# of } u \in C\},$$

where  $C := \{u \in \mathbb{R}^m \mid u^\top B = 0, u \geq 0\}$  is the **projection cone**

## Lemma (Farkas' lemma)

*If  $P = \{x \in \mathbb{R}^d \mid Ax \leq b\}$  is a  $H$ -polyhedron and  $c^\top x \leq \delta$  is valid for  $P$ , then either*

- $P = \emptyset$  or
- $c^\top x \leq \delta'$  is positive combination of  $Ax \leq b$  for some  $\delta' \leq \delta$

## Lemma (Farkas' lemma)

*If  $P = \{x \in \mathbb{R}^d \mid Ax \leq b\}$  is a  $H$ -polyhedron and  $c^\top x \leq \delta$  is valid for  $P$ , then either*

- $P = \emptyset$  or
- $c^\top x \leq \delta'$  is positive combination of  $Ax \leq b$  for some  $\delta' \leq \delta$

*that is, either*

- $\exists u \geq 0 : u^\top A = 0, u^\top b = -1$  or
- $\exists u \geq 0 : u^\top A = c^\top, u^\top b = \delta' \leq \delta$

# Polar of a polytope is an intersection

To prove other direction of the fundamental theorem, we **polarize**

If  $P$  is ( $V$ - or  $H$ -)polytope in  $\mathbb{R}^d$  with  $0 \in \text{int}(P)$ , **polar** of  $P$  is

$$P^\Delta := \{z \in \mathbb{R}^d \mid \forall x \in P : x^\top z \leq 1\}$$

# Polar of a projection is an intersection

To prove other direction of the fundamental theorem, we **polarize**

If  $P$  is ( $V$ - or  $H$ -)polytope in  $\mathbb{R}^d$  with  $0 \in \text{int}(P)$ , **polar** of  $P$  is

$$P^\Delta := \{z \in \mathbb{R}^d \mid \forall x \in P : x^\top z \leq 1\}$$

Then (two last eqs. from Farkas' lemma):

$$(\text{conv}\{v_1, \dots, v_n\})^\Delta = \{x \in \mathbb{R}^d : v_1^\top x \leq 1, \dots, v_n^\top x \leq 1\}$$

$$\left(\{x \in \mathbb{R}^d : A_1 x \leq 1, \dots, A_m x \leq 1\}\right)^\Delta = \text{conv}\{A_1^\top, \dots, A_m^\top\}$$

$$(P^\Delta)^\Delta = P$$



# Polar of a projection is an intersection

To prove other direction of the fundamental theorem, we **polarize**

If  $P$  is ( $V$ - or  $H$ -)polytope in  $\mathbb{R}^d$  with  $0 \in \text{int}(P)$ , **polar** of  $P$  is

$$P^\Delta := \{z \in \mathbb{R}^d \mid \forall x \in P : x^\top z \leq 1\}$$

Then (two last eqs. from Farkas' lemma):

$$(\text{conv}\{v_1, \dots, v_n\})^\Delta = \{x \in \mathbb{R}^d : v_1^\top x \leq 1, \dots, v_n^\top x \leq 1\}$$

$$\left(\{x \in \mathbb{R}^d : A_1 x \leq 1, \dots, A_m x \leq 1\}\right)^\Delta = \text{conv}\{A_1^\top, \dots, A_m^\top\}$$

$$(P^\Delta)^\Delta = P$$

**Exercise.** Prove that the polar of projecting polytope  $Q$  into  $x$ -space is intersecting polytope  $Q^\Delta$  with  $x$ -space, and use this to prove other direction of fundamental theorem.

# Elimination blows up the number of inequalities

If  $Q$  is defined by  $m$  constraints

Then  $\text{proj}_x(Q)$  is defined by at most  $\frac{m^2}{4}$  inequalities  
... after projecting out 1 variable.

# Elimination blows up the number of inequalities

If  $Q$  is defined by  $m$  constraints

Then  $\text{proj}_x(Q)$  is defined by at most  $\frac{m^2}{4}$  inequalities  
... after projecting out 1 variable.

**Exercise.** Verify that this is tight. Find an upper bound when 2 variables are projected out. Is this bound tight?

# Elimination blows up the number of inequalities

If  $Q$  is defined by  $m$  constraints

Then  $\text{proj}_x(Q)$  is defined by at most  $\frac{m^2}{4}$  inequalities  
... after projecting out 1 variable.

**Exercise.** Verify that this is tight. Find an upper bound when 2 variables are projected out. Is this bound tight?

$\implies$  when projecting out  $k$  variables: **exponential blow-up!**

# Elimination blows up the number of inequalities

If  $Q$  is defined by  $m$  constraints

Then  $\text{proj}_x(Q)$  is defined by at most  $\frac{m^2}{4}$  inequalities  
... after projecting out 1 variable.

**Exercise.** Verify that this is tight. Find an upper bound when 2 variables are projected out. Is this bound tight?

$\implies$  when projecting out  $k$  variables: **exponential blow-up!**

*EFs try to exploit this phenomenon: by adding few variables,  
decrease much the number of inequalities*

# Faces of a polytope

## Definition (Face)

If  $P$  is a polytope in  $\mathbb{R}^d$  and  $c^\top x \leq \delta$  is valid for  $P$ , then

$$F := P \cap \{x \in \mathbb{R}^d \mid c^\top x = \delta\} \quad \text{is a \textcolor{red}{face} of } P$$

# Faces of a polytope

## Definition (Face)

If  $P$  is a polytope in  $\mathbb{R}^d$  and  $c^\top x \leq \delta$  is valid for  $P$ , then

$$F := P \cap \{x \in \mathbb{R}^d \mid c^\top x = \delta\} \quad \text{is a \textcolor{red}{face} of } P$$

Particular cases:

- $\emptyset$  is a face, of dimension  $-1$  (use  $0^\top x \leq 1$ )
- a face of dimension 0 is a **vertex**
- a face of dimension 1 is an **edge**
- a face of dimension  $\dim(P) - 1$  is a **facet**
- $P$  is a face of itself, of dimension  $\dim(P)$  (use  $0^\top x \leq 0$ )

# Faces of a polytope

## Definition (Face)

If  $P$  is a polytope in  $\mathbb{R}^d$  and  $c^\top x \leq \delta$  is valid for  $P$ , then

$$F := P \cap \{x \in \mathbb{R}^d \mid c^\top x = \delta\} \quad \text{is a \textcolor{red}{face} of } P$$

Particular cases:

- $\emptyset$  is a face, of dimension  $-1$  (use  $0^\top x \leq 1$ )
- a face of dimension 0 is a **vertex**
- a face of dimension 1 is an **edge**
- a face of dimension  $\dim(P) - 1$  is a **facet**
- $P$  is a face of itself, of dimension  $\dim(P)$  (use  $0^\top x \leq 0$ )

$$\mathcal{L}(P) = (\mathcal{F}(P), \subseteq) \quad \text{\textcolor{red}{face lattice} of } P$$




- 1 A **V-cone** is the nonnegative hull of a finite set of vectors

$$C = \text{cone}\{g_1, \dots, g_k\}$$

- 2 A **H-cone** is the intersection of a finite # of *linear* halfspaces

$$C = \{x \in \mathbb{R}^d \mid Ax \geq 0\}$$




- ① A **V-cone** is the nonnegative hull of a finite set of vectors

$$C = \text{cone}\{g_1, \dots, g_k\}$$

- ② A **H-cone** is the intersection of a finite # of *linear* halfspaces

$$C = \{x \in \mathbb{R}^d \mid Ax \geq 0\}$$



$$\forall C \subseteq \mathbb{R}^d : C \text{ is a } V\text{-cone} \iff C \text{ is a } H\text{-cone}$$


# General polyhedra

- ① A **V-polyhedron** is the sum of a polytope and a cone

$$P = \text{conv}\{v_1, \dots, v_n\} + \text{cone}\{g_1, \dots, g_k\}$$

- ② A **H-polyhedron** is the intersection of a finite # of halfspaces

$$P = \{x \in \mathbb{R}^d \mid Ax \leq b\}$$




# General polyhedra

- ① A **V-polyhedron** is the sum of a polytope and a cone

$$P = \text{conv}\{v_1, \dots, v_n\} + \text{cone}\{g_1, \dots, g_k\}$$

- ② A **H-polyhedron** is the intersection of a finite # of halfspaces

$$P = \{x \in \mathbb{R}^d \mid Ax \leq b\}$$




$\forall P \subseteq \mathbb{R}^d : P \text{ is a V-polyhedron} \iff P \text{ is a H-polyhedron}$

# Today

- 1 Introduction
- 2 Basics on polytopes and polyhedra
- 3 Factorization theorem**

# Slack matrices


... Of polytopes




$$P = \{x \mid A_1 x \leq b_1, \dots, A_m x \leq b_m\}$$

# Slack matrices

... Of polytopes




$$P = \{x \mid A_1 x \leq b_1, \dots, A_m x \leq b_m\}$$




$$P = \text{conv}\{v_1, \dots, v_n\}$$

# Slack matrices

... Of polytopes



$$P = \{x \mid A_1x \leq b_1, \dots, A_mx \leq b_m\}$$





$$P = \text{conv}\{v_1, \dots, v_n\}$$

## Definition

**Slack matrix**  $S \in \mathbb{R}_{+}^{m \times n}$  of polytope  $P$ :

$$S_{ij} := b_i - A_i v_j$$





# Nonnegative factorizations

## Definition

A **rank- $r$  nonnegative factorization** of  $S \in \mathbb{R}^{m \times n}$  is


$$S = TU \quad \text{where} \quad T \in \mathbb{R}_{+}^{m \times r} \quad \text{and} \quad U \in \mathbb{R}_{+}^{r \times n}$$

# Nonnegative factorizations

## Definition

A **rank- $r$  nonnegative factorization** of  $S \in \mathbb{R}^{m \times n}$  is

$$S = TU \quad \text{where} \quad T \in \mathbb{R}_{+}^{m \times r} \quad \text{and} \quad U \in \mathbb{R}_{+}^{r \times n}$$




# Nonnegative factorizations

## Definition

A **rank- $r$  nonnegative factorization** of  $S \in \mathbb{R}^{m \times n}$  is

$$S = TU \quad \text{where} \quad T \in \mathbb{R}_{+}^{m \times r} \quad \text{and} \quad U \in \mathbb{R}_{+}^{r \times n}$$



## Definition (nonnegative rank of $S$ )


$$\text{rk}_{+}(S) := \min\{r \mid \exists \text{ rank-}r \text{ nonnegative factorization of } S\}$$

# Nonnegative factorizations

## Definition

A **rank- $r$  nonnegative factorization** of  $S \in \mathbb{R}^{m \times n}$  is

$$S = TU \quad \text{where} \quad T \in \mathbb{R}_{+}^{m \times r} \quad \text{and} \quad U \in \mathbb{R}_{+}^{r \times n}$$



## Definition (nonnegative rank of $S$ )

$$\begin{aligned} \text{rk}_{+}(S) &:= \min\{r \mid \exists \text{ rank-}r \text{ nonnegative factorization of } S\} \\ &= \min\{r \mid S \text{ is sum of } r \text{ nonnegative rank-1 matrices}\} \end{aligned}$$

# Factorization theorem


## Theorem (Yannakakis' factorization theorem)

*Let  $P$  be a polytope with  $\dim(P) \geq 1$ .*

*For every slack matrix  $S$  of  $P$ :  $xc(P) = rk_+(S)$*

# Factorization theorem

Example: regular polygon



Halfspaces:  $\ell_1^+, \ell_2^+, \dots$

Folding sequence of vertex  $v$ :  $v^{(0)} = v, v^{(1)}, v^{(2)}, \dots$

Folding sequence of facet  $F$ :  $F^{(0)} = F, F^{(1)}, F^{(2)}, \dots$

Slack ( $v^{(i)}$  w.r.t.  $F^{(i)}$ ) = slack ( $v^{(i+1)}$  w.r.t.  $F^{(i+1)}$ ) + **correction**