



Contents lists available at ScienceDirect

Chaos, Solitons and Fractals

Nonlinear Science, and Nonequilibrium and Complex Phenomena

journal homepage: www.elsevier.com/locate/chaos

Frontiers

A quasi-periodic route to chaos in a parametrically driven nonlinear medium



Ana M. Cabanas^{a,*}, Ronald Rivas^{b,c}, Laura M. Pérez^b, Javier A. Vélez^{b,d}, Pablo Díaz^e, Marcel G. Clerc^f, Harald Pleiner^g, David Laroze^{b,*}, Boris A. Malomed^h

^a Sede Esmeralda, Universidad de Tarapacá, Av. Luis Emilio Recabarren 2477, Iquique, Chile

^b Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica, Chile

^c Departamento de Física, Universidad Nacional Experimental Francisco de Miranda, Coro, Venezuela

^d Centro de Nanociencia y Nanotecnología, CEDENNA, Av. Libertador Bernardo O'Higgins 3363, Santiago, Chile

^e Departamento de Ciencias Físicas, Universidad de La Frontera, Casilla 54-D, Temuco, Chile

^f Departamento de Física and Millennium Institute for Research in Optics, Facultad de Ciencias Físicas y Matemáticas, Universidad de Chile, Casilla 4873, Santiago, Chile

^g Max Planck Institute for Polymer Research, D 55021 Mainz, Germany

^h Department of Physical Electronics, School of Electrical Engineering, Faculty of Engineering, Tel Aviv University, Tel Aviv 69978, Israel

ARTICLE INFO

Article history:

Received 2 May 2021

Accepted 20 May 2021

Keywords:

Parametric forcing
Dissipative systems
Pendula chain
Chaos
Lyapunov exponents

ABSTRACT

Small-sized systems exhibit a finite number of routes to chaos. However, in extended systems, not all routes to complex spatiotemporal behavior have been fully explored. Starting from the sine-Gordon model of parametrically driven chain of damped nonlinear oscillators, we investigate a route to spatiotemporal chaos emerging from standing waves. The route from the stationary to the chaotic state proceeds through quasi-periodic dynamics. The standing wave undergoes the onset of oscillatory instability, which subsequently exhibits a different critical frequency, from which the complexity originates. A suitable amplitude equation, valid close to the parametric resonance, makes it possible to produce universal results. The respective phase-space structure and bifurcation diagrams are produced in a numerical form. We characterize the relevant dynamical regimes by means of the largest Lyapunov exponent, the power spectrum, and the evolution of the total intensity of the wave field.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

A natural way to pump energy into various dynamical systems is offered by the use of direct or parametric resonances [1]. Direct resonances were exploited at the dawn of modern physics by Galileo [2] in the characterization of a pendulum. Parametric resonances were highlighted by the pioneering works of Faraday [3] in the study of vibration modes of a water tank. Theoretical and experimental studies of parametrically forced nonlinear systems are relevant to many settings in physics, chemistry, biology, and engineering. Due to the competition between injection and dissipation of energy, the driven systems demonstrate complex behavior, leading to the formation of a wide range of spatiotemporal structures, such as localized ones, extended patterns, and chaotic states, among others [4–11]. One of the most generic scenarios of the self-organizing behavior is pattern formation. With the in-

crease of the energy-injection rate, basic stationary patterns may become unstable and bifurcate into a great variety of more complex ones [6,7,11–22]. In particular, a manifestation of the complex spatiotemporal dynamics is time-aperiodic behavior, as observed, inter alia, in fluids [23–27], chemical waves [28], cardiac fibrillation [29], granular matter [30], electroconvection [31], and photonic devices [32–34]. In the broad variety of physical setups exhibiting complex spatiotemporal behaviors, routes leading to the emergence of spatiotemporal chaos are not fully understood yet, as analytical investigation of the underlying nonlinear partial differential equations is a challenging problem. Diverse routes have been identified for the transition from order to disorder in wave settings, such as defect-mediated turbulence [35–37], onset of spatiotemporal chaos in Galilean-invariant systems [38], spatiotemporal intermittence [39], quasi-periodicity of travelling [40] and standing waves [14], crisis of spatiotemporally chaotic saddles [41–43], spatiotemporal chaos which sets in via stationary branching shocks and holes [44], and the phase turbulence [45].

A generic model that describes periodically forced systems is based on the parametrically driven damped nonlinear Schrödinger

* Corresponding author.

E-mail addresses: ana.cabanas.plana@gmail.com (A.M. Cabanas), dlaroze@uta.cl (D. Laroze).

(PDDNLS) equation [46–48]. This amplitude equation produces a variety of temporal behaviors, including stationary, periodic, and chaotic regimes, such as Faraday waves [3,49–51], single-soliton [47,52–55] and two-soliton [56,57] states, and spatiotemporal chaos [47,58]. Remarkable hydrodynamic modes are excited by the parametric instability in the form of standing (Faraday) waves on the surface of a vertically vibrated Newtonian fluid [3]. These standing waves respond strongest at a half of the forcing frequency (the 2 : 1 resonance) [59]. The basic PDDNLS equation describes parametric instabilities close to the 2 : 1 resonance in weakly dissipative systems [60]. However, this equation in its simplest form is not able to produce certain phenomena, such as stable localized modes that connect a uniform oscillatory state to an extended wave [61], or domain walls between uniform oscillations with opposite phases [62–64]. To study the existence and properties of a broad class of dynamical states, a generalized PDDNLS equation, produced by a systematic expansion procedure, can be used. It adequately accounts for diverse types of the dynamical behavior, such as fronts and confined patterns [65].

This work aims to investigate dynamical scenarios based on standing waves in a parametrically driven chain of pendula, which is considered in the continuum approximation. Starting from the respective parametrically-driven damped sine-Gordon model, it is shown that spatiotemporal chaos can emerge from standing waves. The chaos appears as an extension of quasi-periodic dynamics. Namely, changing parameters of the model, we observe that the standing wave exhibits an oscillatory instability. Further variation of the parameters leads to a secondary oscillatory instability, which eventually gives rise to the spatiotemporal chaos. The generalized PDDNLS equation, valid close to the parametric resonance, allows to systematically investigate the emergence of the complex behavior in a numerical form. The largest Lyapunov exponent, power spectrum, and time evolution of the total norm of the wave field are computed to characterize the transitions between stationary, quasi-periodic, and chaotic dynamical states.

The manuscript is organized as follows. The physical model, the nonlinear analysis and the dynamical indicators used to describe the system are presented in Section 2. Results produced by systematic numerical investigation of the patterns and characterization of the dynamical behavior of the system are reported in Section 3. Finally, conclusions are formulated in Section 4.

2. The parametrically driven damped sine-Gordon equation and the nonlinear-Schrödinger approximation

We start by consideration of a chain of parametrically driven damped pendula (an extended Scott’s model [49]), described in the continuum limit by the parametrically driven damped sine-Gordon equation [50]:

$$\ddot{\theta} = -[\omega_0^2 + \gamma_0 \sin(\omega t)] \sin \theta - \mu_0 \dot{\theta} + \kappa \partial_{xx} \theta, \quad (1)$$

where $\theta(x, t)$ is the angle between the pendulum at position x and the vertical axes at time t , the overdot stands for the time derivative, ω_0 is the eigenfrequency of the pendulum, γ_0 and ω are the amplitude and frequency of the parametric forcing, μ_0 is the damping coefficient and κ the coefficient of the elastic coupling. In the following, by means of rescaling of x and t we fix $\kappa = \omega_0 = 1$. Fig. 1 illustrates different types of dynamical oscillatory states in the system.

The trivial solution $\theta(x, t) = 0$ can be made unstable by the parametric forcing with the frequency close to the resonant (double) value,

$$\omega = 2(\omega_0 + \nu) \quad (2)$$

where ν is a detuning parameter, with $|\nu| \ll \omega_0$. The commonly known analysis [1] produces the stability boundary (the so-called *Arnold tongue*), in the form of

$$\nu^2 + (\mu_0/2)^2 = (\gamma_0/4)^2, \quad (3)$$

depicted in Fig. 2 by the dashed black hyperbola in the parameter plane of the detuning and forcing strength. While crossing




Fig. 1. Spatiotemporal dynamical behavior produced by the parametrically driven damped sine-Gordon equation (1). Color-coded spatiotemporal distributions of the continuous field, $\theta(x, t)$, are displayed, with time interval $\Delta\tau = 2000$, in the middle row, parallel to typical configurations of the chain of coupled pendula shown in the top row, which is approximated by Eq. (1) in the continuum limit. a) A spatially uniform state obtained at the amplitude and detuning of the parametric drive $\gamma_0 = 0.85$ and $\nu = -0.02$ (see Eq. (2)), with the largest Lyapunov exponent $\lambda_{\max} = -0.037$. b) A regular spatially periodic pattern at $\gamma_0 = 0.40$ and $\nu = -0.05$, with $\lambda_{\max} = -0.041$. c) A chaotic state at $\gamma_0 = 0.61$ and $\nu = -0.08$, with $\lambda_{\max} = 0.055$. Bottom panels d), e), and f) show normalized Fourier spectra $S(f)$ and phase portraits in the plane of $(\theta, \dot{\theta})$ at a fixed spatial point ($x = 0$), plotted for a), b) and c), respectively. Other parameters in Eq. (1) are $\omega_0 = 1, \mu_0 = 0.1$, and $\kappa = 1$.




Fig. 2. The diagram of dynamical states in the (γ, ν) parameter plane of Eqs. (1) and (2). The dashed curve represents the instability boundary (Arnold tongue) given by Eq. (3). Green circles (●) represent spatially uniform states, red triangles (▼) denote standing waves, blue squares (■) chaotic waves, and black diamonds (◆) correspond to quasi-periodic states with small amplitudes. Here and in Fig. 5 below, the trivial state, $\theta = 0$, occurs in the white area. Right panels display snapshots of the respective spatial patterns. The fixed parameters are the same as in Fig. 1. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

the boundary, the destabilization occurs through a supercritical (subcritical) bifurcation at positive (negative) values of the detuning [60], uniform oscillations taking place inside of the “tongue” [64]. The corresponding power spectrum, displayed in panel d) of Fig. 1 shows well-defined peaks determined by the forcing. The respective inset shows the phase portrait corresponding to a closed limit cycle.

In the Arnold tongue, the uniform oscillations are observed at small detuning, as illustrated in Fig. 2 by green circles. Making the detuning more negative, the uniform oscillations develop a supercritical oscillatory spatial instability with a nonzero wavenumber [6]. It is characterized by emergence of a novel frequency, which is not commensurate with the frequency of the uniform oscillations. The instability gives rise to standing waves, as shown in panels b) and e) of Fig. 1. Note that in the respective phase portrait the limit cycle transforms into a torus. These standing waves are denoted by red triangles in the diagram of states displayed in Fig. 2. Further decreasing the detuning, the standing wave develops a complex spatiotemporal behavior, as illustrated in panels c) and e) of Fig. 1. It is seen that the temporal spectrum at a fixed spatial point now represents dynamical chaos, and, likewise, the phase portrait corresponds to a strange attractor [8]. Such a complex spatiotemporal behavior, denoted by blue squares in Fig. 2, is observed in a broad region of the Arnold tongue.

To characterize the dynamical nature of the complex waves, we use the largest Lyapunov exponent λ_{\max} [66]. First, $\lambda_{\max} < 0$ corresponds to a stable equilibrium state, a uniform or patterned one. On the contrary, $\lambda_{\max} > 0$ implies chaotic dynamics, with exponential sensibility to initial conditions. The case of $\lambda_{\max} = 0$ corresponds to quasi-periodic behavior or complex behavior with non-exponential sensitivity. Fig. 3 shows λ_{\max} as a function of detuning ν (see Eq. (2)) for different values of the drive’s strength, $\gamma_0 = \{0.4, 0.5, 0.6, 0.7\}$. The plots make it possible to distinguish regular and chaotic behavior, which, as said above, correspond to $\lambda_{\max} < 0$ and $\lambda_{\max} > 0$, respectively.

Thus, the parametrically driven damped sine-Gordon equation (1) provides the transition from the spatially uniform oscillations to the spatiotemporal chaos, via the intermediate spatially periodic pattern. To achieve a better grasp of these dynamical regimes, in the next section we develop analysis under conditions of weak




Fig. 3. The largest Lyapunov exponent λ_{\max} vs. detuning ν (see Eq. (2)), at different values of the strength of the parametric forcing: $\gamma_0 = \{0.4, 0.5, 0.6, 0.7\}$. The other parameters are the same as in Fig. 2.

nonlinearity and slowly-varying amplitude, close to the parametric resonance.

2.1. The weak-nonlinearity analysis

The above model, based in the sine-Gordon equation, can be essentially simplified for small-amplitude solutions, whose main harmonic frequency is close to the eigenfrequency, ω_0 . To this end, the following ansatz is adopted [60]:

$$\theta(x, t) = \psi(\tau, X)e^{-i\omega\tau/2} + \text{c.c.} + \text{h.o.t.}, \quad (4)$$

where $\psi(\tau, X)$ is a slowly complex envelope amplitude, $\tau \equiv t/4$, $X \equiv x/\sqrt{2}$, while c.c. and h.o.t. stand for the complex conjugate and higher-order terms, respectively. Close to the 2 : 1 parametric resonance, the frequency is $\omega = 2(1 + \nu)$, as per Eq. (2). The substitution of ansatz (4) in (1) yields, in the first approximation, the PDDNLS equation [47], which is the basic model of the driven systems close to the 2 : 1 parametric resonance:

$$\frac{\partial \psi}{\partial \tau} = -i\nu\psi - i|\psi|^2\psi - i\frac{\partial^2 \psi}{\partial X^2} - \mu\psi + \gamma\bar{\psi}, \quad (5)$$

where $\mu \equiv \mu_0/2$, $\gamma \equiv \gamma_0/4$, and $\bar{\psi}$ is the complex conjugate of ψ . This equation is known to produce stationary, time-periodic, or chaotic solutions, including Faraday waves [50,62], soliton-like modes [47,52], two-soliton [56,57] and soliton-antisoliton [68] bound states, and spatiotemporal chaos [47,58].

The PDDNLS equation admits different spatially uniform solutions. The instability boundary of the trivial one, $\psi = 0$, is

$$\gamma^2 = \mu^2 + \nu^2, \quad (6)$$

cf. Eq. (3). In addition, the trivial solution suffers an instability which sets in at $\gamma = \mu$ for the positive detuning [50,67]. Nontrivial uniform solutions of Eq. (5) are

$$\psi_{\pm, \pm} = \pm x_0(1 \pm iy_0), \quad (7)$$

where $x_0 = \sqrt{(\gamma - \mu)(\phi - \nu)/2\gamma}$ and $y_0 = \sqrt{(\mu - \gamma)/(\mu + \gamma)}$ with $\phi \equiv \sqrt{\gamma^2 - \mu^2}$, and two independent signs \pm . Note that the uniform solutions emerge at $\gamma = \mu$ through a saddle-node bifurcation. They are unstable in the parameter space of Eq. (5), with the exception of the case of zero detuning ($\nu = 0$), in which case $\psi_{\pm, +}$ is marginally stable [63].

In fact, the underlying sine-Gordon equation (1) may have stable solutions for uniform oscillations. To provide this possibility, as well as the existence of more complex states that bifurcate from the uniform oscillations, it is necessary to proceed to the generalized PDDNLS equation. Such a generalization can be written as

$$\frac{\partial \psi}{\partial \tau} = -i\nu\psi - i|\psi|^2\psi - i\frac{\partial^2 \psi}{\partial X^2} - \mu\psi + \gamma\bar{\psi} + \mathcal{N}_\psi, \quad (8)$$




Fig. 4. The spatiotemporal evolution of the real part of ψ in established dynamical states shown within a time interval $\Delta\tau = 2000$ (the first and second rows), and snapshots of the respective spatial patterns at a certain time (the third row). a), b) and c): Two regular small-amplitude patterns $\psi_{+/-}$ (in- and out-of-phase ones, see the text) at $\gamma = 0.85$ and $\nu = -0.25$, with $\lambda_{\max+} = -1.853 \times 10^{-3}$ and $\lambda_{\max-} = -1.852 \times 10^{-3}$. d), e) and f): Quasi-periodic states $\psi_{+/-}$ at $\gamma = 0.85$ and $\nu = -0.45$, with $\lambda_{\max+} = 4.92 \times 10^{-4}$ and $\lambda_{\max-} = 5.13 \times 10^{-4}$. g), h) and i): Small-amplitude chaotic patterns $\psi_{+/-}$ at $\gamma = 0.85$ and $\nu = -0.55$, with $\lambda_{\max+} = 1.288 \times 10^{-2}$ and $\lambda_{\max-} = 1.359 \times 10^{-2}$. Fixed parameters are $b = 1/12$, $\delta = 4/15$, $\beta = -1/24$, $\alpha = -0.65$, $a = 1/6$ and $\mu = 0.275$.




Fig. 5. Phase diagrams in the (γ, ν) parameter plane for different initial conditions shown in the insets. a): A flat profile with a bump; b) a spatially-periodic profile. Green circles (●) correspond to the spatially uniform state, red triangles (▼) denote stationary small-amplitude states, blue squares (■) the small-amplitude chaotic pattern, and black diamonds (◆) correspond to quasi-periodic states with a small amplitude. c) Coincidence regions, in which the different initial conditions, used in diagrams a) and b), produce identical final states. The fixed parameters are the same as in Fig. 4. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

with additional terms \mathcal{N}_ψ produced by the weak-nonlinearity expansion of underlying equations of the sine-Gordon type [60,64]. Thus, at order $\gamma^{5/2}$ of the expansion, one can derive, in the general case,

$$\mathcal{N}_\psi = \gamma(b\bar{\psi}|\psi|^4 + \delta\psi^3 + \beta\psi^3|\psi|^2) + ia\psi|\psi|^4 + \eta \frac{\partial^2 \psi}{\partial X^2} + c\psi|\psi|^2 + i\zeta\bar{\psi} \left(\frac{\partial \psi}{\partial X} \right)^2, \quad (9)$$

where η accounts for the diffusion, c represents the nonlinear dissipation, (b, δ, β) are coefficients of the nonlinear parametric forcing, a controls the quintic nonlinearity, and ζ is the coefficient of the nonlinear-drift term. The amended equation (8) has been employed to produce kinks, localized patterns and traveling pulses [51,60–65,68–76]. In fact, not all the additional terms are necessary to secure the stability of the uniform states inside of the Arnold tongue [71]. Note that, in the case of the sine-Gordon equation, the expansion at order $\gamma^{5/2}$ yields $\eta = 0$, $c = 0$, $\alpha = 1/2$, $\delta = -1/6$, $a = 1/6$, $b = 1/12$, $\beta = -1/24$, and $\zeta = 0$ [65].

We here focus on the case of $\eta = 0$ and $c = 0$, as effects of the terms $\sim \eta$ and c have been studied before [71–73]. Besides, the effect of ζ is neglected, as it is known too, inducing traveling solutions [74]. Then, it is easy to find that the squared amplitude of the uniform-state solution, $|\psi|^2$, is determined by equation

$$\gamma^2 = \frac{\mu^2}{[1 + (\delta + \alpha)|\psi|^2 + (\beta + b)|\psi|^4]^2}$$

$$+ \frac{(\nu + |\psi|^2 - a|\psi|^4)^2}{[1 + (\alpha - \delta)|\psi|^2 + (b - \beta)|\psi|^4]^2}. \quad (10)$$

To study small-amplitude patterns generated by the extended PDDNLS equation, the coefficients of \mathcal{N}_ψ in Eq. (9) are introduced as free parameters. In particular, numerical results are displayed below for $b = 1/12$, $\delta = 4/15$, $\beta = -1/24$, $\alpha = -0.65$, $a = 1/6$, and $\mu = 0.275$, which make it possible to produce generic findings.

2.2. Indicators of complexity and transition between dynamical regimes

To characterize the dynamics produced by Eq. (8), we use standard indicators, one of which is the time-dependent energy. It is often used to study non-regular dynamics in fluids, optics [75–80] and other physical systems [81–83]:

$$Q(\tau) = \frac{1}{2L} \int_{-L}^{+L} |\psi(\tau, X)|^2 dX, \quad (11)$$

where $2L$ is the system's size. The simple uniform stationary regime has $Q = \text{const}$, and it is a (quasi-)periodic function of time in the case of (quasi-)periodic dynamics. An apparently random time dependence of $Q(\tau)$ corresponds to chaotic behavior.

To categorize different dynamical regimes, we also use the power spectrum of the amplitude, $S_Q(f) = |\mathfrak{F}(f)|^2$, where \mathfrak{F} is the Fourier transform of $Q(\tau)$,

$$\mathfrak{F}(f) = \frac{1}{\sqrt{2\pi}} \int_0^{\tau_{\max}} Q(\tau) \exp(-if\tau) d\tau. \quad (12)$$

If the time series is regular, function $S_Q(f)$ has a finite number of quasi-discrete peaks. On the contrary, the spectrum is essentially continuous if the series is chaotic [84]. In addition, to characterize spatial fluctuations of field ψ , we also use the time average of the local density,

$$\mathcal{L}(x) = \frac{1}{T - T_0} \int_{T_0}^T |\psi(\tau, X)|^2 d\tau, \quad (13)$$

and its spatial power spectrum,

$$S_{\mathcal{L}}(k) = \int_{-L}^{+L} S_{\mathcal{L}}(k) \exp(-ikX) dX, \quad (14)$$

at a set of points in the frequency space $k = (k_1, \dots, k_n)$. Similar to the temporal power spectrum, its spatial counterpart, $S_{\mathcal{L}}(k)$, helps to quantify different types of the spatial behavior of the system.

Further, a well-known potent indicator of the dynamics is the largest Lyapunov exponent [66,85–106]

$$\lambda_{\max} = \lim_{\tau \rightarrow \infty} \left[\frac{1}{\tau} \ln \left(\frac{\|\delta\psi(\tau, x)\|}{\|\delta\psi_0\|} \right) \right], \quad (15)$$

where $\|\bullet\|$ stands for the quadratic norm, and $\delta\psi(\tau, x)$ is a numerically generated solution of the linearized equation, $\partial(\delta\psi)/\partial\tau = \mathbf{J} \cdot \delta\psi$, with \mathbf{J} being the Jacobian matrix of Eq. (8). We remark that λ_{\max} predicts how fast distance $\delta\psi$ between two initially close trajectories of field ψ grows in the course of the evolution. Namely, if the system is chaotic with $\lambda_{\max} > 0$, two configurations, that were close at $\tau = \tau_0$, separate in phase space exponentially fast. On the other hand, in the case of $\lambda_{\max} < 0$ they converge to a stationary attractor. The marginal case, $\lambda_{\max} = 0$, corresponds to time-periodic, quasi-periodic, or complex dynamics without exponential sensibility to the variation of initial conditions.

3. The quasi-periodic route to spatiotemporal chaos in the generalized PDDNLS equation

3.1. Numerically obtained patterns

A variable-step fifth-order Runge-Kutta scheme [107] has been used for simulations of Eq. (8) with the precision of 10^{-7} . The spatial derivatives were approximated by means of the second-order central finite-difference method, and the Neumann's boundary conditions were used. Typical the half-size $L = 100$ and $N = 1500$ lattice points imply that the mesh size of the numerical scheme is $\Delta x = 200/1500 \approx 0.13$. Sensitivity of the findings to the discretization has been verified by changing N and Δx . Variation of the domain size L was used to verify that finite-size effects do not affect the results. We have used two essentially different initial conditions, as shown below. The integration was performed over sufficiently long times (up to $\tau = 4.8 \times 10^3$, to check that transients have faded out.

For the analysis of spatial patterns we present in Fig. 4 the spatiotemporal evolution of the real part of the wave field, $\psi_R \equiv \text{Re}\{\psi(x, \tau)\}$. Starting close to the tip of the Arnold tongue, which implies the consideration of a small detuning ($\gamma = 0.85$) and the forcing strength slightly greater than the dissipation ($\nu = -0.25$, see Eq. (6)). The regular in-phase and out-of-phase states (ψ_{\pm} , with $\psi_R > 0$ and $\psi_R < 0$, respectively) are observed in Figs. 4a-c. In these panels the spatial periodicity observed corresponds to the real part of $\exp(ikx)$ with some k , while $|\psi|$ remains constant. The values of λ_{\max} for these regular patterns are negative: $\lambda_{\max+} = -1.853 \times 10^{-3}$ and $\lambda_{\max-} = -1.852 \times 10^{-3}$, respectively.

Following the decrease of the detuning and/or the increase of the forcing strength, the pattern exhibits an oscillatory instability, which gives rise to standing waves. Figs. 4d-f show the corresponding temporarily quasi-periodic states at $\gamma = 0.85$ and $\nu = -0.45$,

with $\lambda_{\max+} = 4.92 \times 10^{-4}$ and $\lambda_{\max-} = 5.13 \times 10^{-4}$ (actually, these values may be considered as a numerical zero). Changing the parameters further in the same direction, the standing wave exhibits an oscillatory instability, producing complex spatiotemporal patterns. Figs. 4 g-i) display typical examples of the chaotic spatiotemporal evolution of the complex patterns, accounted for by $\lambda_{\max} > 0$.

Fig. 5 shows phase diagrams inside the Arnold tongue for two different initial conditions (see insets): a) a flat profile with a bump, and b) a small-amplitude periodic pattern. The two respective diagrams are similar with a few differences, see the comparison between them in panel c) which displays regions in which both initial conditions produce the same final state. In the case of the small-amplitude spatially periodic initial condition, the region of the uniform state (green circles) is somewhat smaller, and the distribution of regions of temporarily stationary (red triangles), quasi-periodic (black diamonds), and chaotic (blue squares) patterns is shifted to smaller values of detuning ν , while quasi-periodic states appear at larger values of γ . In both cases, the dynamical chaos occupies the upper left corner, while near to the zero-detuning line, $\nu = 0$, the uniform state is the dominant one. Thus, the small-amplitude chaos is equally likely to be generated by both initial conditions.

3.2. Dynamics inside the Arnold tongue

Panel a) of Fig. 6 presents the dependence of λ_{\max} on the forcing strength in the case of a spatially periodic pattern chosen as initial condition. In this plot, we observe the transition from a stationary solution (uniform or patterned one) to an oscillatory one (the standing wave), which then turns into a spatiotemporal chaotic pattern, following the increase of the forcing strength, γ . The growth of λ_{\max} with the subsequent increase of γ indicates that the chaotic spatiotemporal pattern becomes more complex. Furthermore, around $\gamma \approx 2.2$, we observe an abrupt transition from $\lambda_{\max} > 0$ to small islands with $\lambda_{\max} = 0$, which corresponds to a phenomenon known as the *crisis of a strange attractor* [84].

Further, the patterns are characterized by the plot for $Q_{\max}(\gamma)$ as a function of the forcing strength γ . Fig. 6 b) displays it at $\nu = 0.1$. It is produced by repeatedly picking the maximum value of the energy function $Q(\tau)$, see Eq. (11), from temporal intervals after the disappearance of transient features in the simulated evolution. The system is stationary or periodic if there is a unique value of Q_{\max} , while a continuous distribution of Q_{\max} in a finite interval implies temporal quasi-periodicity or chaos. This figure shows that energy Q strongly depends on forcing γ , showing several transitions between regular and chaotic states, denoted by the same symbols as in Fig. 5. The transitions are shown in greater detail in the inset, where the plot is blown up by a factor of 5.4. The diagrams for λ_{\max} and Q_{\max} provide mutually complimentary descriptions of the oscillatory, quasi-periodic, and chaotic behavior. Note that the plot of Q_{\max} is represented by a thin curve at $\gamma < 1.8$, becoming fuzzy at larger values of γ . In particular, alternation of islands, from chaos to regularity, is observed in the range of $1.92 < \gamma < 2.0$. The system becomes chaotic again at $\gamma > 2.0$ through an abrupt transition, and remains chaotic at larger values of γ considered here.

To grasp the route to chaos in a clearer form, we have computed the spatial Fourier spectrum $S_L(k)$, as per Eq. (14). Fig. 6 shows $S_L(k)$ for the spatially uniform state at $\gamma = 1.50$ in panel c), for a temporally quasi-periodic one at $\gamma = 1.95$ in d), and for a chaotic state at $\gamma = 2.1$ in e). Increase of the number of frequency peaks in the spectrum is observed as γ increases at a fixed value of the detuning, $\nu = -0.1$. These plots corroborate that Eq. (8) produces a standing wave with a well-defined wavenumber




Fig. 6. Characterization of the route to spatiotemporal chaos produced by simulations of Eq. (8), is shown by means of dynamical indicators. a) λ_{\max} , for $\nu = -0.1$, and b) Q_{\max} , for $\nu = +0.1$, as a function of the forcing strength, γ , see Eqs. (11) and (15). The insets are zoomed-in versions of the main plots. The meaning of the symbols is the same as in Fig. 5: green circles correspond to spatially uniform states, red triangles to stationary small-amplitude ones, and blue squares to the small-amplitude chaotic pattern. Panel c) shows the spatial Fourier spectrum $S_L(k)$ (see Eq. (14)) for a regular state at $\gamma = 1.50$, d) a temporarily quasi-periodic state at $\gamma = 1.95$, and e) a chaotic one at $\gamma = 2.1$. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 7. The power spectrum $S_Q(f)$ of the norm (energy) $Q(t)$, defined as per Eq. (11). a) A quasi-periodic state at $\gamma = 1.89$ and $\nu = -0.1$. b) A chaotic state at $\gamma = 0.85$ and $\nu = -0.55$. The insets show the respective plots of $Q(t)$.

and its harmonics in Fig. 6 c). As the forcing strength increases, the system generates incommensurate wavenumbers, which causes the spectral peaks to spread, see Fig. 6 d). Further increase of the strength makes the former sharp peaks broad in Fig. 6 e), which is a hallmark of spatial chaos.

Further understanding of these patterns is obtained by considering the temporal evolution of the energy function $Q(t)$, defined by Eq. (11), and the corresponding Fourier power spectrum, $S_Q(f)$. Panel a) of Fig. 7 shows $S_Q(f)$ for a quasi-periodic pattern at $\gamma = 1.89$ and $\nu = -0.1$, with $\lambda_{\max} = -3.234 \times 10^{-4}$. The spectrum features a set of discrete peaks at particular frequencies. Accordingly, the inset shows regular evolution of $Q(t)$. On the other hand, panel b) of Fig. 7 exhibits a chaotic state at $\gamma = 0.85$ and $\nu = -0.55$, with $\lambda_{\max} = 1.288 \times 10^{-2}$ showing a continuous spectrum $S_Q(f)$, typical for chaotic states, with irregular evolution of $Q(t)$ in the respective inset. Thus, the transition to the spatiotemporal chaos proceeds through an intermediate stage which features the temporal quasi-periodicity.

4. Conclusions

In this work, we have investigated the transition to spatiotemporal chaos from small-amplitude patterns in damped driven systems close to the parametric resonance. Starting from the continuum limit of a chain of driven pendula, which is based on the equation of the sine-Gordon type, the analysis is performed in

the framework of the generalized PDDNLS (parametrically driven damped nonlinear Schrödinger) equation, which is produced by the small-amplitude expansion of the underlying sine-Gordon equation. The systematic numerical analysis demonstrates that the onset of the spatiotemporal chaos proceeds through an intermediate temporally-quasi-periodic route. We use the generalized PDDNLS equation (8), which contains higher-order nonlinear terms, because the standard PDDNLS model, Eq. (5), fails to produce stable spatially uniform solutions, which play a crucial role in obtaining localized states. In particular, we have found the small-amplitude patterns that are stationary, quasi-periodic, or chaotic in time depending on the values of the strength of the parametric forcing, γ , and resonance detuning, ν . We have concentrated on chaotic solutions, using different tools to characterize their dynamical behavior, such as the time dependence of the norm (energy) Q , the corresponding power spectrum, and the maximum Lyapunov exponent. Small-amplitude chaotic patterns exist inside of the Arnold tongue, competing with regular ones. Varying γ at constant ν or vice versa leads in various transitions between these states. There are well confined chaotic patterns and others, in which the chaotic area penetrates the homogeneous one, following the increase of γ .

The universal nature of the present model suggests that similar scenarios of the transition to the small-amplitude spatiotemporal chaos may be expected in physical settings such as vertically oscillating fluid layers, magnetic systems, forced nonlinear lattices, and

optical waveguides. A promising direction for the extension of this work is to extend it to the two-dimensional case.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

LMP and DL appreciate the hospitality of the MPI-P (Mainz) during their stay in Germany. LMP, PD and DL acknowledge partial financial support from FONDECYT 1180905. MGC acknowledges partial financial support from Millennium Institute for Research in Optics, ANID–Millennium Science Initiative Program–ICN17_012 and FONDECYT 1180903. JAV and DL acknowledge partial financial support from Centers of excellence with BASAL/CONICYT financing, Grant AFB180001, CEDENNA. The work of BAM is supported, in part, by the Israel Science Foundation through grant No. 1286/17. This author also acknowledges support from Instituto de Alta Investigación, Universidad de Tarapacá (Arica, Chile).

References

- [1] Landau LD, Lifshitz EM. *Mechanics*. Pergamon Press; 1976.
- [2] Prova A, Marenzana M. *Thus Spoke Galileo: The Great Scientist's Ideas and their Relevance to the Present Day*. Oxford University Press; 2006.
- [3] Faraday M. On a peculiar class of acoustical figures; and on certain forms assumed by groups of particles upon vibrating elastic surfaces. *Philos Trans R Soc London* 1831;121:299.
- [4] Nocolis G, Prigogine I. *Self-Organization in Nonequilibrium Systems*. New York: J Wiley and Sons; 1977.
- [5] Haken H. *Synergetics*. Physics Bulletin 1977;28:412.
- [6] Cross MC, Hohenberg PC. Pattern formation outside of equilibrium. *Rev Mod Phys* 1993;65:851.
- [7] Pismen LM. *Patterns and interfaces in dissipative dynamics*. Berlin: Springer; 2006.
- [8] Manneville P. *Instabilities, chaos and turbulence*. London: Imperial college Press; 2004.
- [9] Walgraef D. *Spatio-temporal pattern formation: with examples from physics, chemistry, and materials science*. Springer Science & Business Media; 2012.
- [10] Murray JD. *Mathematical Biology*. Berlin: Springer-Verlag; 1993.
- [11] Nocolis G. *Introduction to Nonlinear Science*. Cambridge: Cambridge University Press; 1995.
- [12] Couillet P, Gil L, Lega J. Defect-mediated turbulence. *Phys Rev Lett* 1989a;62:1619.
- [13] Couillet P, Gil L, Lega J. A form of turbulence associated with defects. *Physica D* 1989b;37:91.
- [14] Clerc MG, Verschuere N. Quasiperiodicity route to spatiotemporal chaos in one-dimensional pattern-forming systems. *Phys Rev E* 2013;88:052916.
- [15] Brand HR, Deissler RJ. Interaction of localized solutions for subcritical bifurcations. *Phys Rev Lett* 1989;63:2801.
- [16] Brand HR, Deissler RJ. Stable spatially localized solutions and holes in optical bistability. *Physica A* 1994;204:87.
- [17] Deissler RJ, Brand HR. Periodic, quasiperiodic, and chaotic localized solutions of the quintic complex Ginzburg-Landau equation. *Phys Rev Lett* 1994;72:478.
- [18] Deissler RJ, Brand HR. Two-dimensional localized solutions for subcritical bifurcations in systems with broken rotational symmetry. *Phys Rev E* 1995a;51:R852.
- [19] Deissler RJ, Brand HR. Interaction of Breathing Localized Solutions for Subcritical Bifurcations. *Phys Rev Lett* 1995b;74:4847.
- [20] Cartes C, Descalzi O. Collisions of non-explosive dissipative solitons can induce explosions. *Physical Review A* 2016;93:031801.
- [21] Descalzi O, Brand HR. Collisions of non-explosive dissipative solitons can induce explosions. *Chaos* 2018;28:075508.
- [22] Descalzi O, Brand HR. *Physical Review E* 2020;101:040201.
- [23] Decker W, Pesch W, Weber A. Spiral defect chaos in Rayleigh-Bénard convection. *Phys Rev Lett* 1994;73:648.
- [24] Echebarria B, Riecke H. Defect Chaos of Oscillating Hexagons in Rotating Convection. *Phys Rev Lett* 2000;84:4838.
- [25] Daniels KE, Bodenschatz E. Defect turbulence in inclined layer convection. *Phys Rev Lett* 2002;88:034501.
- [26] Miranda MA, Burguete J. Experimentally observed route to spatiotemporal chaos in an extended one-dimensional array of convective oscillators. *Phys Rev E* 2009;79:046201.
- [27] Brunet P, Limat L. Defects and spatiotemporal disorder in a pattern of falling liquid columns. *Phys Rev E* 2004;70:046207.
- [28] Ouyang Q, Flesselles JM. Transition from spirals to defect turbulence driven by a convective instability. *Nature (London)* 1996;379:143.
- [29] Qu Z, Weiss JN, Garfinkel A. Spatiotemporal Chaos in a Simulated Ring of Cardiac Cells. *Phys Rev Lett* 1997;78:1387.
- [30] Moon SJ, Shattuck MD, Bizon C, Goldman DI, Swift JB, Swinney HL. Phase diagram of vertically shaken granular matter. *Phys Rev E* 2001;65:011301.
- [31] Zhou SQ, Ahlers G. Spatiotemporal chaos in electroconvection of a homeotropically aligned nematic liquid crystal. *Phys Rev E* 2006;74:046212.
- [32] Verschuere N, Bortolozzo U, Clerc MG, Residori S. Spatiotemporal Chaotic Localized State in Liquid Crystal Light Valve Experiments with Optical Feedback. *Phys Rev Lett* 2013;110:104101.
- [33] Clerc MG, Gonzalez-Cortes G, Odent V, Wilson M. Optical textures: characterizing spatiotemporal chaos. *Opt Express* 2016;24:15478.
- [34] Clerc MG, Alvarez-Garrido F, Gonzalez-Cortes G. Transition to Spatiotemporal Intermittency and Defect Turbulence in Systems under Translational Coupling. *Phys Rev Lett* 2020;124:164101.
- [35] Couillet P, Lega J. Defect-Mediated Turbulence in Wave Patterns. *Europhys Lett* 1988;7:511.
- [36] Couillet P, Gil L, Lega J. Defect-mediated turbulence. *Phys Rev Lett* 1989c;62:1619.
- [37] Couillet P, Gil L, Lega J. A form of turbulence associated with defects. *Physica D* 1989d;37:91.
- [38] Goren G, Eckmann JP, Procaccia I. Scenario for the onset of space-time chaos. *Phys Rev E* 1998;57:4106.
- [39] Chate H. Spatiotemporal intermittency regimes of the one-dimensional complex Ginzburg-Landau equation. *Nonlinearity* 1994;7:185.
- [40] Brochard F, Gravier E, Bonhomme G. Transitions to spatiotemporal chaos and turbulence of flute instabilities in a low - β magnetized plasma. *Phys Rev E* 2006;73:036403.
- [41] He K. From Quantum Dynamics to the Canonical Distribution: General Picture and a Rigorous Example. *Phys Rev Lett* 1998;80:696.
- [42] He K, Chian ACL. Critical dynamic events at the crisis of transition to spatiotemporal chaos. *Phys Rev E* 2004;69:026207.
- [43] Rempel EL, Chian ACL. Origin of Transient and Intermittent Dynamics in Spatiotemporal Chaotic Systems. *Phys Rev Lett* 2007;98:014101.
- [44] Sherratt JA, Smith MJ. Transition to spatiotemporal chaos via stationary branching shocks and holes. *Physica D* 2012;241:1671.
- [45] Kuramoto Y. *Chemical Oscillations, Waves, and Turbulence*. New York: Springer; 1984.
- [46] Miles JW. Parametrically excited solitary waves. *J Fluid Mech* 1984;148:451.
- [47] Barashenkov IV, Bogdan MM, Korobov VI. Stability Diagram of the Phase-Locked Solitons in the Parametrically Driven, Damped Nonlinear Schrödinger Equation. *Europhys Lett* 1991;15:113.
- [48] Malomed BA. Nonlinearity and discreteness: Solitons in lattices. In: Kevrekidis PG, Cuevas-Maraver J, Saxena A, editors. *Emerging Frontiers in Nonlinear Science*, pp. 81-110. Springer Nature Switzerland AG: Cham; 2020.
- [49] Scott AC. A Nonlinear Klein-Gordon Equation. *Am J Phys* 1969;37:52-61.
- [50] Couillet P, Firsch T, Sonnino G. Dispersion-induced patterns. *Phys Rev E* 1994;49:2087.
- [51] Clerc MG, Coulibaly S, Laroze D. Interaction law of 2D localized precession states. *EPL* 2010a;90:38005.
- [52] Barashenkov IV, Zemlyanaya EV, van Heerden TC. Time-periodic solitons in a damped-driven nonlinear Schrödinger equation. *Phys Rev E* 2011;83:056609.
- [53] Alexeeva NV, Barashenkov IV, Tsironis GP. Impurity-Induced Stabilization of Solitons in Arrays of Parametrically Driven Nonlinear Oscillators. *Phys Rev Lett* 2000;84:3053.
- [54] Barashenkov IV, Zemlyanaya EV. Stable Complexes of Parametrically Driven, Damped Nonlinear Schrödinger Solitons. *Phys Rev Lett* 1999;83:2568.
- [55] Zemlyanaya EV, Alexeeva NV. Oscillating solitons of the driven, damped nonlinear Schrödinger equation. *Theoretical and Mathematical Physics* 2009;159:870.
- [56] Barashenkov IV, Zemlyanaya EV. Soliton complexity in the damped-driven nonlinear Schrödinger equation: Stationary to periodic to quasiperiodic complexes. *Phys Rev E* 2011;83:056610.
- [57] Urzagasti D, Laroze D, Clerc MG, Coulibaly S, Pleiner H. Two-soliton precession state in a parametrically driven magnetic wire. *J Appl Phys* 2012;111:07D111.
- [58] Shchesnovich VS, Barashenkov IV. Soliton-radiation coupling in the parametrically driven, damped nonlinear Schrödinger equation. *Physica D* 2002;164:83.
- [59] Arnold VI. *Geometrical methods in the theory of ordinary differential equations*. New York: Springer-Verlag; 1983.
- [60] Clerc MG, Coulibaly S, Laroze D. Localized chaotic patterns in weakly dissipative systems. *Int J Bif Chaos* 2009a;19:3525.
- [61] Clerc MG, Coulibaly S, Laroze D. Effective-parametric resonance in a non-oscillating system. *EPL* 2012;97:30006.
- [62] Clerc MG, Coulibaly S, Laroze D. Localized states and non-variational Ising-Bloch transition of a parametrically driven easy-plane ferromagnetic wire. *Physica D* 2010b;1239:72.
- [63] Clerc MG, Coulibaly S, Laroze D. Localized states beyond the asymptotic parametrically driven amplitude equation. *Phys Rev E* 2008;77:056209.
- [64] Clerc MG, Coulibaly S, Laroze D. Nonvariational Ising-Bloch Transition in Parametrically Driven Systems. *Int J BifChaos* 2009b;19:2717.
- [65] Urzagasti D, Laroze D, Pleiner H. Localized chaotic patterns in weakly dissipative systems. *J Special Topics* 2014a;223:141.

- [66] Pikovsky A, Politi A. *Lyapunov Exponents: A Tool to Explore Complex Dynamics*. Cambridge: Cambridge University Press; 2016.
- [67] Elphick C, Tirapegui E, Brachet ME, Coulet P, Iooss G. A simple global characterization for normal forms of singular vector fields. *Physica D* 1987;29:95.
- [68] Urzagasti D, Aramayo A, Laroze D. Soliton-antisoliton interaction in a parametrically driven easy-plane magnetic wire. *Phys Lett A* 2014b;378:2614.
- [69] Urzagasti D, Laroze D, Pleiner H. Two-dimensional localized chaotic patterns in parametrically driven systems. *Phys Rev E* 2017;95:052216.
- [70] Kenig E, Malomed BA, Cross MC, Lifshitz R. Intrinsic localized modes in parametrically driven arrays of nonlinear resonators. *Phys Rev E* 2009;80:046202.
- [71] Barashenkov IV, Cross S, Malomed BA. Multistable pulselike solutions in a parametrically driven Ginzburg-Landau equation. *Phys Rev E* 2003;68:056605.
- [72] Burke J, Yochelis A, Knobloch E. Classification of Spatially Localized Oscillations in Periodically Forced Dissipative Systems. *SIAM J Appl Dyn Sys* 2008;7:651.
- [73] Ma Y-P, Burke J, Knobloch E. Defect-mediated snaking: A new growth mechanism for localized structures. *Physica D* 2010;239:1867.
- [74] León AO, Clerc MG, Coulibaly S. Traveling pulse on a periodic background in parametrically driven systems. *Phys Rev E* 2015;91: 050901(R)
- [75] Edri Y, Meron E, Yochelis A. Spatial asymmetries of resonant oscillations in periodically forced heterogeneous media. *Physica D* 2020a;410:132501.
- [76] Edri Y, Meron E, Yochelis A. Spatial heterogeneity may form an inverse camel shaped Arnold's tongue in parametrically forced oscillations. *Chaos* 2020b;30:023120.
- [77] Akhmediev N, Soto-Crespo JM, Town G. Pulsating solitons, chaotic solitons, period doubling, and pulse coexistence in mode-locked lasers: Complex Ginzburg-Landau equation approach. *Phys Rev E* 2001;63:056602.
- [78] Ankiewicz A, Devine N, Akhmediev N, Soto-Crespo JM. Continuously self-focusing and continuously self-defocusing two-dimensional beams in dissipative media. *Phys Rev A* 2008;77:033840.
- [79] Akhmediev N, Soto-Crespo JM. Exploding solitons and Shil'nikov's Theorem. *Phys Lett A* 2003;317:287.
- [80] Akhmediev N, Soto-Crespo JM. Strongly asymmetric soliton explosions. *Phys Rev E* 2004;70:036613.
- [81] Batiste O, Knobloch E, Alonso A, Mercader I. Spatially localized binary-fluid convection. *J Fluid Mech* 2006;560:149.
- [82] Burke J, Knobloch E. Localized states in the generalized Swift-Hohenberg equation. *Phys Rev E* 2006;73:056211.
- [83] Burke J, Knobloch E. Homoclinic snaking: Structure and stability. *Chaos* 2007;17:037102.
- [84] Ott E. *Chaos in Dynamical Systems*. Cambridge: Cambridge University Press; 1993.
- [85] Wolf A, Swift JB, Swinney HL, Vastano JA. Determining Lyapunov exponents from a time series. *Physica D* 1985;16:285.
- [86] Sprott JC. *Chaos and Time-Series Analysis*. Oxford: Oxford: Oxford University Press; 2003.
- [87] Laroze D, Bragard J, Suarez OJ, Pleiner H. Characterization of the Chaotic Magnetic Particle Dynamics. *IEEE Trans Mag* 2011;47:3032.
- [88] Bragard J, Pleiner H, Suarez OJ, Vargas P, Gallas JAC, Laroze D. Chaotic dynamics of a magnetic nanoparticle. *Phys Rev E* 2011;84:037202.
- [89] Laroze D, Becerra-Alonso D, Gallas JAC, Pleiner H. Magnetization Dynamics Under a Quasiperiodic Magnetic Field. *IEEE Trans Mag* 2012;48:3567.
- [90] Eckmann JP, Ruelle D, Ciliberto S. Lyapunov exponents from time series. *Phys Rev A* 1986;34:4971.
- [91] Geist K, Parlitz U, Lauterborn W. Comparison of different methods for computing Lyapunov exponents. *Prog Theor Phys* 1990;83:875.
- [92] Scheel JD, Cross MC. Lyapunov exponents for small aspect ratio Rayleigh-Bénard convection. *Phys Rev E* 2006;74:066301.
- [93] Karimi A, Paul MR. Quantifying spatiotemporal chaos in Rayleigh-Bénard convection. *Phys Rev E* 2012;85:046201.
- [94] Bonatto C, Gallas JAC. Periodicity Hub and Nested Spirals in the Phase Diagram of a Simple Resistive Circuit. *Phys Rev Lett* 2008;101:054101.
- [95] Ramirez-Avila GM, Gallas JAC. How similar is the performance of the cubic and the piecewise-linear circuits of Chua? *Phys Lett A* 2010;375:143.
- [96] Gallas JAC. The Structure of Infinite Periodic and Chaotic Hub Cascades in Phase Diagrams of Simple Autonomous Flows. *Thermal convection in a nonlinear non-Newtonian magnetic fluid*. *Int J Bif Chaos* 2010;20:197.
- [97] Vélez JA, Bragard J, Pérez LM, Cabanas AM, Suarez OJ, Laroze D, Mancini HL. Periodicity characterization of the nonlinear magnetization dynamics. *Chaos* 2020;30:093112.
- [98] Mahmud MN, Siri Z, Vélez JA, Pérez LM, Laroze D. Chaotic convection in an Oldroyd viscoelastic fluid in saturated porous medium with feedback control. *Chaos* 2020;30:073109.
- [99] Laroze D, Pleiner H. Thermal convection in a nonlinear non-Newtonian magnetic fluid. *Commun Nonlinear Sci Numer Simulat* 2015;26:167.
- [100] Laroze D, Siddheshwar PG, Pleiner H. Chaotic convection in a ferrofluid. *Commun Nonlinear Sci Numer Simulat* 2013;18:2436.
- [101] Cabanas AM, Clerc MG, Laroze D, Leon AO. Chaotic patterns and localized states in spin valves. *J Magn Magn Mater* 2019;476:589.
- [102] Cabanas AM, Pérez LM, Laroze D. Strange non-chaotic attractors in spin valve systems. *J Magn Magn Mater* 2018;460:320.
- [103] Pérez LM, Bragard J, Mancini HL, Gallas JAC, Cabanas AM, Suarez OJ, Laroze D. Effect of anisotropies on the magnetization dynamics. *Netw Heterog Media* 2015;10:209.
- [104] Brugnago EL, Gallas JAC, Beims MW. Predicting regime changes and durations in Lorenz's atmospheric convection model. *Chaos* 2020;30:083106.
- [105] Cabanas AM, Vélez JA, Pérez LM, Dáz P, Clerc MG, Laroze D, Malomed BA. Dissipative structures in a parametrically driven dissipative lattice: Chimera, localized disorder, continuous-wave, and staggered states. *Chaos, Solitons and Fractals* 2021;146:110880.
- [106] Siddheshwar PG, Kanchana C, Laroze D. A study of Darcy-Bénard regular and chaotic convection using a new local thermal non-equilibrium formulation. *Physics of Fluids* 2021;33:044107.
- [107] Press WH, Teukolsky SA, Vetterling WT, Flannery BP. *Numerical Recipes in FORTRAN*. UK: Cambridge University Press; 1992.